

Non-linear Simulations of ELMs in ASDEX Upgrade: Status and Plans

Matthias Hölzl, Isabel Krebs,
Karl Lackner, Sibylle Günter

- 1 JOREK
- 2 Localized ELMs
- 3 Mode Interaction
- 4 Filaments
- 5 Field Penetration
- 6 Plans
- 7 Summary

- 1 JOREK
- 2 Localized ELMs
- 3 Mode Interaction
- 4 Filaments
- 5 Field Penetration
- 6 Plans
- 7 Summary

- ▶ Originally developed at CEA Cadarache
[G. Huysmans and O. Czarny. *Nucl.Fusion*, 47, 659 (2007)]
- ▶ Non-linear reduced MHD in toroidal geometry (next slide)
- ▶ Two-fluid extensions
- ▶ Full MHD in development

- ▶ 2D Bezier finite elements + Fourier decomposition
- ▶ Fully implicit time evolution
- ▶ GMRES with physics-based preconditioning
- ▶ MPI + OpenMP parallelization

$$\frac{\partial \Psi}{\partial t} = \eta j - R [\mathbf{u}, \Psi] - F_0 \frac{\partial \mathbf{u}}{\partial \phi}$$

$$\frac{\partial \rho}{\partial t} = -\nabla \cdot (\rho \mathbf{v}) + \nabla \cdot (D_{\perp} \nabla_{\perp} \rho) + S_{\rho}$$

$$\frac{\partial (\rho T)}{\partial t} = -\mathbf{v} \cdot \nabla (\rho T) - \gamma \rho T \nabla \cdot \mathbf{v} + \nabla \cdot (K_{\perp} \nabla_{\perp} T + K_{\parallel} \nabla_{\parallel} T) + S_T$$

$$\mathbf{e}_{\phi} \cdot \nabla \times \left\{ \rho \frac{\partial \mathbf{v}}{\partial t} = -\rho (\mathbf{v} \cdot \nabla) \mathbf{v} - \nabla p + \mathbf{j} \times \mathbf{B} + \mu \Delta \mathbf{v} \right\}$$

$$\mathbf{B} \cdot \left\{ \rho \frac{\partial \mathbf{v}}{\partial t} = -\rho (\mathbf{v} \cdot \nabla) \mathbf{v} - \nabla p + \mathbf{j} \times \mathbf{B} + \mu \Delta \mathbf{v} \right\}$$


$$\mathbf{j} \equiv -\mathbf{j}_{\phi} = \Delta^* \Psi$$

$$\boldsymbol{\omega} \equiv -\boldsymbol{\omega}_{\phi} = \nabla_{\text{pol}}^2 \mathbf{u}$$


Variables: $\Psi, \mathbf{u}, \mathbf{j}, \boldsymbol{\omega}, \rho, T, v_{\parallel}$

Definitions: $\mathbf{B} \equiv \frac{F_0}{R} \mathbf{e}_{\phi} + \frac{1}{R} \nabla \Psi \times \mathbf{e}_{\phi}$ and $\mathbf{v} \equiv -R \nabla \mathbf{u} \times \mathbf{e}_{\phi} + v_{\parallel} \mathbf{B}$


[H. R. Strauss. *Phys.Fluids*, 19, 134 (1976)]



- ▶ Input from CLISTE
- ▶ Initial grid (Grids shown with reduced resolution)
- ▶ Flux aligned grid including X-point(s)
- ▶ Equilibrium flows
- ▶ Time-integration




- ▶ Input from CLISTE
- ▶ Initial grid (Grids shown with reduced resolution)
- ▶ Flux aligned grid including X-point(s)
- ▶ Equilibrium flows
- ▶ Time-integration



- ▶ Input from CLISTE
- ▶ Initial grid (Grids shown with reduced resolution)
- ▶ Flux aligned grid including X-point(s)
- ▶ Equilibrium flows
- ▶ Time-integration

- ▷ **Matthias Hölzl** (PostDoc)
 - Edge localized modes
 - Magnetic perturbations
 - Resistive wall model
- ▷ **Isabel Krebs** (PhD)
 - JOREK for master thesis and first part of her PhD
 - Mode interaction during ELMs
 - Going to Princeton in September
 - Benchmark with M3D-C1 planned
- ▷ **Alexander Lessig** (PhD starting in June)
 - Different ELM types
 - Full ELM cycles
 - Comparisons to experiments
- ▷ **Nina Wenke** (Practical till July)
 - CLISTE-JOREK interface
 - Visualization

- ▷ Typical ASDEX Upgrade H-mode equilibrium
- ▷ Resistivity too large by factor 10 due to numerical constraints (improving)




$$T_e + T_i \approx 12 \text{ keV}$$

$$n_e \approx 6 \cdot 10^{19} \text{ m}^{-3}$$


- 1 JOREK
- 2 Localized ELMs
- 3 Mode Interaction
- 4 Filaments
- 5 Field Penetration
- 6 Plans
- 7 Summary

- ASDEX Upgrade: Expanded and localized ELMs observed (distribution)




Signature of a Solitary Magnetic Perturbation in ASDEX Upgrade

[R. P. Wenninger, et al. *Nucl.Fusion*, 42, 114025 (2012)]



$$n = 0, 8, 16$$

- ▶ Red/blue surfaces correspond to 70 percent of maximum/minimum values




$$n = 0, 1, 2, 3, 4, \dots, 16$$

- ▶ Red/blue surfaces correspond to 70 percent of maximum/minimum values
- ▶ Localized due to several strong harmonics with adjacent n



⇒ **Similar to Solitary Magnetic Perturbations**

[M. Hölzl, et al. *38th EPS*, P2.078 (2011); M. Hölzl, et al. *Phys. Plasmas*, 19, 082505 (2012b)]

- 1 JOREK
- 2 Localized ELMs
- 3 Mode Interaction**
- 4 Filaments
- 5 Field Penetration
- 6 Plans
- 7 Summary




Example for ELM signature with strong low-n component



Histogram of dominant components in a TCV discharge (23 ELMs)

[R. P. Wenninger, et al. Nucl.Fusion (submitted)]




- ▶ Non-linear drive of low-n modes
- ▶ Start with simplified case including $n = 0, 4, 8, 12, 16$ (periodicity 4)

- ▶ Quadratic terms lead to mode coupling $(n_1, n_2) \leftrightarrow n_1 \pm n_2$
- ▶ For instance: $(16, 12) \leftrightarrow 4$
- ▶ Model assuming mode rigidity and fixed background:

$$\begin{aligned}
 \dot{A}_4 &= \overbrace{\gamma_4 A_4}^{\text{linear}} + \overbrace{\gamma_{8,-4} A_8 A_4 + \gamma_{12,-8} A_{12} A_8 + \gamma_{16,-12} A_{16} A_{12}}^{\text{non-linear interaction}} \\
 \dot{A}_8 &= \gamma_8 A_8 + \gamma_{4,4} A_4 A_4 + \gamma_{12,-4} A_{12} A_4 + \gamma_{16,-8} A_{16} A_8 \\
 \dot{A}_{12} &= \gamma_{12} A_{12} + \gamma_{4,8} A_4 A_8 + \gamma_{16,-4} A_{16} A_4 \\
 \dot{A}_{16} &= \gamma_{16} A_{16} + \gamma_{8,8} A_8 A_8 + \gamma_{4,12} A_4 A_{12}
 \end{aligned}$$


- ▶ Linear growth rates from JOREK simulation + Energy conservation
- ▶ Determine few free parameters by minimizing quadratic differences

[I. Krebs. Master's thesis, LMU, Munich (2012)]




- ▷ Non-linear drive recovered
- ▷ Saturation not recovered (of course)

Mode Interaction Simple Model (3)



- ▶ Applied to full simulation with $n = 0 \dots 16$


[I. Krebs, et al. *Phys. Plasmas* (submitted)]




- ▶ Applied to full simulation with $n = 0 \dots 16$
- ▶ **Explains low- n features in experimental observations**

[I. Krebs, et al. *Phys. Plasmas* (submitted)]


- 1 JOREK
- 2 Localized ELMs
- 3 Mode Interaction
- 4 Filaments**
- 5 Field Penetration
- 6 Plans
- 7 Summary




- ▷ Energy time traces during an ELM crash
- ▷ Simulation with $n = 0,8$




- ▷ Energy time traces during an ELM crash
- ▷ Simulation with $n = 0,8$




Detached filaments quickly lose their pressure due to fast parallel heat conduction. Substructures appear in divertor heat flux patterns.




Detached filaments quickly lose their pressure due to fast parallel heat conduction. Substructures appear in divertor heat flux patterns.




Detached filaments quickly lose their pressure due to fast parallel heat conduction. Substructures appear in divertor heat flux patterns.




Detached filaments quickly lose their pressure due to fast parallel heat conduction. Substructures appear in divertor heat flux patterns.




Detached filaments quickly lose their pressure due to fast parallel heat conduction. Substructures appear in divertor heat flux patterns.




Detached filaments quickly lose their pressure due to fast parallel heat conduction. Substructures appear in divertor heat flux patterns.




Detached filaments quickly lose their pressure due to fast parallel heat conduction. Substructures appear in divertor heat flux patterns.




Detached filaments quickly lose their pressure due to fast parallel heat conduction. Substructures appear in divertor heat flux patterns.



Detached filaments quickly lose their pressure due to fast parallel heat conduction. Substructures appear in divertor heat flux patterns.




Detached filaments quickly lose their pressure due to fast parallel heat conduction. Substructures appear in divertor heat flux patterns.




Detached filaments quickly lose their pressure due to fast parallel heat conduction. Substructures appear in divertor heat flux patterns.

- 1 JOREK
- 2 Localized ELMs
- 3 Mode Interaction
- 4 Filaments
- 5 Field Penetration**
- 6 Plans
- 7 Summary



Pseudocolor
Var: abs_curr
6.000
4.500
3.000
1.500
0.000
Max: 5.897
Min: 0.000


- ▶ Equilibrium with $q = 1$ surface:
Influence on plasma rotation
(W. Suttrop, M. Dunne)
- ▶ Highly preliminary (weekend...)
- ▶ ELM-Coils in ASDEX Upgrade
- ▶ $n = 1$ configuration
- ▶ Boundary condition from modified
Vacfield
- ▶ Perturbation gradually switched on
- ▶ Differences between resonant and
non-resonant?



Pseudocolor
Var: abs_curr
6.000
4.500
3.000
1.500
0.000
Max: 5.897
Min: 0.000



- ▶ Equilibrium with $q = 1$ surface:
Influence on plasma rotation
(W. Suttrop, M. Dunne)
- ▶ Highly preliminary (weekend...)
- ▶ ELM-Coils in ASDEX Upgrade
- ▶ $n = 1$ configuration
- ▶ Boundary condition from modified
Vacfield
- ▶ Perturbation gradually switched on
- ▶ Differences between resonant and
non-resonant?

- 1 JOREK
- 2 Localized ELMs
- 3 Mode Interaction
- 4 Filaments
- 5 Field Penetration
- 6 Plans**
- 7 Summary



- ▶ ELM types
- ▶ Full ELM cycle
- ▶ Experiment comparisons

With A. Lessig, AUG-Team




16 perturbation coils are currently installed in ASDEX Upgrade

[W. Suttrup, et al. *24th IAEA, EX/3-4* (2012)]

- ▶ Simulate ELM mitigation with magnetic perturbations

With F. Orain, AUG-Team




Discretization of first ITER wall in the STARWALL code which describes vacuum region and wall currents

[P. Merkel and M. Sempf. 21st IAEA, TH/P3-8 (2006); E. Strumberger, et al. 38th EPS, P5.082 (2011)]

- ▶ Interaction of instabilities with conducting structures (RWMs, VDEs, disruptions, . . .) [M. Hölzl, et al. *JPCS*, 401, 012010 (2012a)]

With P. Merkel, G. Huysmans, E. Nardon



First benchmarks done: VDE in ITER-like limiter plasma

- ▷ Interaction of instabilities with conducting structures (RWMs, VDEs, disruptions, . . .) [M. Hölzl, et al. *JPCS*, 401, 012010 (2012a)]



With P. Merkel, G. Huysmans, E. Nardon

- ▶ Issues with code quality
- Significantly improved, non-regression testing established (CEA and IPP)
- ▶ Issues with positivity in “violent” instabilities
 - ▶ Issues with convergence with many toroidal harmonics
 - ▶ Issues with memory consumption and scaling
- Ongoing work (E. Franck NMPP and CEA)

- 1 JOREK
- 2 Localized ELMs
- 3 Mode Interaction
- 4 Filaments
- 5 Field Penetration
- 6 Plans
- 7 Summary**



- **Localized ELMs**
- **Non-linear mode interaction**
- **Filament formation**
- **Field penetration**

- ELM types, full ELM cycle, comparisons
- ELM mitigation
- Resistive walls




- **Localized ELMs**
- **Non-linear mode interaction**
- **Filament formation**
- **Field penetration**

- ELM types, full ELM cycle, comparisons
- ELM mitigation
- Resistive walls




- **Localized ELMs**
- **Non-linear mode interaction**
- **Filament formation**
- **Field penetration**



- ELM types, full ELM cycle, comparisons
- ELM mitigation
- Resistive walls



- **Localized ELMs**
 - **Non-linear mode interaction**
 - **Filament formation**
 - **Field penetration**
-
- ELM types, full ELM cycle, comparisons
 - ELM mitigation
 - Resistive walls



- **Localized ELMs**
 - **Non-linear mode interaction**
 - **Filament formation**
 - **Field penetration**
-
- ELM types, full ELM cycle, comparisons
 - ELM mitigation
 - Resistive walls



References

- M. Hölzl, et al. *38th EPS*, P2.078 (2011).
- M. Hölzl, et al. *JPCS*, 401, 012010 (2012a).
- M. Hölzl, et al. *Phys.Plasmas*, 19, 082505 (2012b).
- G. Huysmans and O. Czarny. *Nucl.Fusion*, 47, 659 (2007).
- I. Krebs. Master's thesis, LMU, Munich (2012).
- I. Krebs, et al. *Phys.Plasmas* (submitted).
- P. Merkel and M. Sempf. *21st IAEA*, TH/P3-8 (2006).
- H. R. Strauss. *Phys.Fluids*, 19, 134 (1976).
- E. Strumberger, et al. *38th EPS*, P5.082 (2011).
- W. Suttrop, et al. *24th IAEA*, EX/3-4 (2012).
- R. P. Wenninger, et al. *Nucl.Fusion*, 42, 114025 (2012).
- R. P. Wenninger, et al. *Nucl.Fusion* (submitted).

Slides and Publications

<http://me.steindaube.de>

Acknowledgements

I. Krebs, K. Lackner, S. Günter,
G. Huysmans, F. Orain, P. Merkel,
E. Nardon, R. Wenninger, E. Strumberger,
M. Dunne, E. Viezzer, M. Bécoulet,
N. Wenke, AUG-Team

$n = 1$ mode structure

