

**Investigations on Sawtooth Reconnection in  
ASDEX Upgrade Tokamak Discharges  
Using the 3D Non-linear Two-fluid MHD Code M3D-C<sup>1</sup>**

I. Krebs<sup>1</sup>, S.C. Jardin<sup>2</sup>, V. Igochine<sup>3</sup>, S. Günter<sup>3</sup>, M. Hoelzl<sup>3</sup> and the ASDEX Upgrade team<sup>3</sup>

<sup>1</sup> *Max-Planck/Princeton Center for Plasma Physics*

<sup>2</sup> *Princeton Plasma Physics Laboratory, Princeton, NJ, USA*

<sup>3</sup> *Max Planck Institute for Plasma Physics, Garching, Germany*

We study sawtooth reconnection in ASDEX Upgrade tokamak [Herrmann et al., Fusion Sci. Technol. 44(3) (2003)] plasmas by means of 3D non-linear two-fluid MHD simulations in toroidal geometry using the high-order finite element code M3D-C1 [S.C. Jardin et al., J. Phys.: Conf. Ser. 125 (2008)]. Parameters and equilibrium of the simulations are based on typical sawtooth ASDEX Upgrade discharges. The simulation results are compared to features of the experimental observations such as the sawtooth crash time and frequency, the evolution of the safety factor profile and the 3D evolution of the temperature. 2D ECE imaging measurements during sawtooth crashes in ASDEX Upgrade indicate that the heat is transported out of the core through a narrow poloidally localized region [Igochine et al., Phys. Plasmas 17 (2010)]. We investigate if incomplete sawtooth reconnection can be seen in the simulations which is suggested by soft X-ray tomography measurements in ASDEX Upgrade showing that an  $(m=1, n=1)$  perturbation is typically observed to survive the sawtooth crash and approximately maintain its radial position [Igochine et al., Phys. Plasmas 17 (2010)].